

06 JUN 2005

RECEIVED

01 MAR 2004

WIPO

PCT

PL 1126982

# THE UNITED STATES OF AMERICA

**TO ALL TO WHOM THESE PRESENTS SHALL COME:**

UNITED STATES DEPARTMENT OF COMMERCE  
United States Patent and Trademark Office

February 25, 2004

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE.

APPLICATION NUMBER: 60/431,267

FILING DATE: December 06, 2002

RELATED PCT APPLICATION NUMBER: PCT/US03/38672

By Authority of the  
COMMISSIONER OF PATENTS AND TRADEMARKS

*M. Tarver*

M. TARVER  
Certifying Officer



**PRIORITY  
DOCUMENT**

SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH RULE 17.1(a) OR (b)

BEST AVAILABLE COPY

12-09-02 12:06/02 60/431267

12/06/02  
10370 U.S. PTO

10370 U.S. PTO  
60/431267

Approved for use through 10/31/2002 OMB 0851-0032  
Patent and Trademark Office, U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number

# **PROVISIONAL APPLICATION FOR PATENT COVER SHEET**

This is a request for filing a PROVISIONAL APPLICATION FOR PATENT under 37 CFR 1.53 (c).

Express Mail Label No. ET008281151US

INVENTOR(S)					
Given Name (first and middle [if any])		Family Name or Surname		Residence (City and either State or Foreign Country)	
Mamix L.		Bosch		Medina, WA	
<input type="checkbox"/> Additional inventors are being named on the separately numbered sheets attached hereto					
TITLE OF THE INVENTION (500 characters max)					
ADMINISTRATION OF DENDRITIC CELLS PARTIALLY MATURED IN VITRO FOR THE TREATMENT OF TUMORS					
CORRESPONDENCE ADDRESS					
Direct all correspondence to:					
<input checked="" type="checkbox"/> Customer Number		20350		<div style="border: 1px solid black; padding: 5px;">Place Customer Number Bar Code Label here</div>	
OR Type Customer Number here					
<input type="checkbox"/> Firm or Individual Name					
Address					
Address					
City		State		ZIP	
Country		Telephone		Fax	
ENCLOSED APPLICATION PARTS (check all that apply)					
<input checked="" type="checkbox"/> Specification Number of Pages		18		<input type="checkbox"/> CD(s), Number	
<input type="checkbox"/> Drawing(s) Number of Sheets				<input type="checkbox"/> Other (specify)	
<input type="checkbox"/> Application Data Sheet. See 37 CFR 1.76					
METHOD OF PAYMENT OF FILING FEES FOR THIS PROVISIONAL APPLICATION FOR PATENT					
<input checked="" type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27.					
<input type="checkbox"/> A check or money order is enclosed to cover the filing fees					
<input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge filing fees or credit any overpayment to Deposit Account Number:				20-1430	
<input type="checkbox"/> Payment by credit card. Form PTO-2038 is attached.				80	
The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.					
<input checked="" type="checkbox"/> No.					
<input type="checkbox"/> Yes, the name of the U.S. Government agency and the Government contract number are:					

Respectfully submitted,  
SIGNATURE Brian W. Poor  
TYPED or PRINTED NAME Brian W. Poor  
TELEPHONE 206-467-9600

Date 12/6/02  
REGISTRATION NO. 32,928  
(if appropriate)  
Docket Number: 020093-004000US

## **USE ONLY FOR FILING A PROVISIONAL APPLICATION FOR PATENT**

This collection of information is required by 37 CFR 1.51. The information is used by the public to file (and by the PTO to process) a provisional application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.14. This collection is estimated to take 8 hours to complete, including gathering, preparing, and submitting the complete provisional application to the PTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, Washington, D.C., 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Box Provisional Application, Assistant Commissioner for Patents, Washington, D.C. 20231. SE 5014968 v1

Attorney Docket No.: 020093-004000

**PROVISIONAL**

**PATENT APPLICATION**

**ADMINISTRATION OF DENDRITIC CELLS PARTIALLY MATURED  
*IN VITRO* FOR THE TREATMENT OF TUMORS**

**Inventor(s):** Marnix L. Bosch, a citizen of the Netherlands  
residing at 7814 N.E. 14th Street, Medina, WA 98039

**Assignee:** Northwest Biotherapeutics, Inc.  
21720 23<sup>rd</sup> Drive SE, Suite 100  
Bothell, Washington 98021

**Entity:** Small business concern

**ADMINISTRATION OF DENDRITIC CELLS PARTIALLY MATURED  
IN VITRO FOR THE TREATMENT OF TUMORS**

**BACKGROUND OF THE INVENTION**

Dendritic cells (DCs) are recognized as the vehicle of choice for active immunotherapy of cancer. Animal experiments have demonstrated the potential of DC based immunotherapy in both protecting mice from tumor formation and eliminating established tumors. These successes have been at least partially duplicated in humans, in small clinical trials. The transition from small safety- or proof-of-concept trials to larger trials in which activity or efficacy can be demonstrated has been hindered by the laborious and cumbersome nature of DC preparation (see below). As a consequence, few companies have been interested in developing DC-based cancer vaccines despite the large potential upside of such products.

Intratumoral (IT) injection of DCs is a special form of DC-based immunotherapy. Upon injection, the DCs take up antigen from apoptotic or dying tumor cells, and present the antigen to T cells after migration to the lymph nodes. Indeed it was found that the efficacy of such treatments in animal models correlates with the degree of apoptosis in the tumor (Candido *et al.*, *Cancer Res.* 61:228-236, 2001), which suggests that this approach is fully compatible with treating tumors with chemotherapeutic agents or radiation prior to the injection of DCs. In addition, several groups have demonstrated that such combination therapy is particularly effective against established tumors (Nikitina *et al.*, *Int. J. Cancer* 94:825-833, 2001; Tanaka *et al.*, *Int. J. Cancer* 101:265-269, 2002; Tong *et al.*, *Cancer Res.* 61:7530-7535, 2001).

Since the tumor cells are the source of antigen, IT injection foregoes the need for both the selection and manufacturing of tumor antigens as they are currently used in most DC based therapy approaches. Selection of a tumor antigen is often driven by the need for companies to have a proprietary position and the few tumor antigens identified to date have not been proven to provide significant clinical benefit. In addition, the use of such tumor antigens often results in a monovalent vaccine, which can lose its effectiveness if the tumor cells down regulate the expression of the antigen used in immunization. Of course, the need

to manufacture the tumor antigen under conditions required under Good Manufacturing Practices (GMP) adds additional cost to classical DC-based immunizations.

IT injection of DCs subjects the dendritic cells to an immunosuppressive tumor environment. Tumors are known to produce cytokines that inactivate the DCs or that have the ability to skew T cell response towards a Th2-type response. Several groups have used genetic modification of DCs to attempt to overcome these suppressive effects, especially through the production of the cytokine Interleukin 12 (IL-12; Nishioka *et al.*, *Cancer Res.* 59:4035-4041, 1999; Melero *et al.*, *Gene Therapy* 6:1779-1784, 1999) or expression of CD40 ligand (Kikuchi *et al.*, *Blood* 96:91-99, 2000). The encouraging results described by these groups further demonstrate the viability of IT injection of DCs as a therapeutic approach.

Triozi *et al.* (*Cancer* 89:2647-2654, 2000) describe IT injection of DCs in patients with metastatic melanoma or breast cancer. They obtained tumor regression in 4 patients with melanoma and in two patients with breast carcinoma. Biopsies of the regressing lesions demonstrated infiltrating T cells, suggesting that the DC had indeed activated an immune response against the tumor cells. Overall these data demonstrated that IT injection of DCs was feasible in humans, and could provide significant clinical benefit. However, significant down regulation of MHC class II antigens and of the B7-2 costimulatory molecule on injected DCs has been observed. Down regulation of these critical molecules would be expected to reduce the immunostimulatory potential of the DCs, but as provided in the present invention this can be avoided through partial maturation of the DCs prior to administration.

DCs exist in peripheral tissues in an immature form, ready to take up and process antigen. It is this immature cell that is most closely mimicked by the DCs generated from monocytes in the presence of GM-CSF and IL-4. Various stimuli can initiate the maturation of DCs, during which process the cells lose their capacity to take up antigen efficiently, and gain their T cell stimulatory functions. This process is complex and at least *in vitro* can take up to 48 hours to complete. One other consequence of maturation is a change in the migratory properties of the cells. For example, maturation induces several chemokine receptors, including CCR7, which direct the cells to the T cell regions of draining lymph nodes, where the mature DCs activate T cells against the antigens that are presented on the DC surface in the context of class I and class II MHC molecules.

The induction of tolerance has been linked to the cross-presentation of (self) antigens by DCs (Kurts *et al.*, *J. Exp. Med.* 186:239-245), and the current prevailing hypothesis posits that immature DC continually present antigens to the immune system to

maintain tolerance (Kurts *et al.*, *J. Exp. Med.* 184:923-930, 1996), suggesting that maturation of DCs *in vivo* is required for efficient immunostimulation. It has now become apparent that DC maturation can result in either immunostimulatory or immunosuppressive DCs (Chakraborty *et al.*, *Clin. Immunol.* 94:88-98, 1999). The precise nature of the maturation stimuli that produce either outcome has not been elucidated, but it is generally accepted that immunostimulatory DCs produce IL-12 and express CD40 ligand (Albert *et al.*, *Nature Immunol.* 2:988-989, 2001; Lanzavecchia, *Haematologica* 84 Suppl. EHA 4:23-25, 1999). Therefore, partially matured DC should be used for IT injection, especially if the stimuli used for maturation of immature dendritic cells are known to result in expression of CD40 ligand and in the production of IL-12.

#### BRIEF SUMMARY OF THE INVENTION

The present invention provides a method for producing an anti-tumor immune response comprising the administration of a cell population comprising dendritic cells that have been partially matured *in vitro* such that the partially matured dendritic cells retain the ability to take up tumor antigen and can induce an immune response subsequent to administration to an individual. The method provides for the efficient up take and processing of tumor antigens within the tumor and tumor bed in spite of the overall immunosuppressive environment within the tumor. This immunosuppressive environment would be expected to interfere with dendritic cell function by negatively impacting on their immunostimulatory properties. Further, mature dendritic cells would not be administered to tumors because the cells do not efficiently process antigen.

Administration of the partially matured dendritic cells can be directly into an existing tumor. Further, administration can be into the tumor bed, the tissue region in and around a tumor either before or after treatment, *i.e.*, surgical removal or resection of the tumor, prior to, simultaneous with, or subsequent to chemotherapy, radiation therapy, or combinations thereof, and the like. The partially matured dendritic cells can also be administered to a lymph node or lymphatic region that directly drains the tumor or surrounding tumor region. In addition the partially matured dendritic cells can be administered into a circulatory vessel or into a lymphatic region that delivers blood or lymph directly into the tumor or tumor bed, or the organ afflicted by or with the tumor. Still further, the partially matured dendritic cells can be administered generally into the circulatory or lymphatic system such that the cells will be delivered into the tumor region.

Dendritic cells or precursors thereof, useful in the present invention can be obtained from, for example, skin, spleen, bone marrow, thymus, lymph nodes, umbilical cord blood, or peripheral blood, and the like. Further, the dendritic cells can be obtained from an individual to be treated or from a healthy individual HLA-matched to the individual to be treated. The dendritic cells or precursors thereof can be cryopreserved prior to being contacted with a maturation agent and formulated into a composition for administration to an individual.

The dendritic cells used in the methods of the invention are partially matured *in vitro*. Suitable agents for the induction of dendritic cell maturation include, for example, but are not limited to, Bacillus Calmette-Guerin (BCG), interferon  $\gamma$  (IFN $\gamma$ ), lipopolysaccharide (LPS), tumor necrosis factor  $\alpha$  (TNF $\alpha$ ), or a combination thereof, and the like. BCG as used in the present invention can comprise whole BCG, cell wall constituents of BCG, BCG-derived lipoarabidomannans, or any other BCG components. In certain embodiments, the BCG is heat-inactivated BCG or formalin-treated BCG. Typically, combinations of BCG and IFN $\gamma$  are used for the partial maturation of the dendritic cells, although as an alternative BCG can be used alone. An effective amount of BCG is typically about  $10^5$  to  $10^7$  cfu per milliliter of tissue culture media and an effective amount of IFN $\gamma$  is typically about 100 to about 1000 Units per milliliter of tissue culture media.

The present invention also comprises compositions comprising dendritic cells partially matured *in vitro* in the presence of a dendritic cell maturation agent combined with a physiologically acceptable carrier, buffer and/or excipient for administration to an individual with a tumor. Partially mature dendritic cells used in the compositions of the invention typically do not express the cell surface marker CD83 that is a characteristic of mature dendritic cells. The composition typically will comprise about  $10^2$  to about  $10^{10}$  partially matured dendritic cells. Subsequent to partial maturation the dendritic cells can be cryopreserved prior to administration to an individual. The partially matured dendritic cells of the compositions can be isolated from the patient having the tumor or the partially matured dendritic cells can have been HLA matched to the patient.

## DETAILED DESCRIPTION OF THE INVENTION

Dendritic cells are a diverse population of antigen presenting cells found in a variety of lymphoid and non-lymphoid tissues. (See Liu, *Cell* 106:259-62 (2001); Steinman, *Ann. Rev. Immunol.* 9:271-96 (1991)). Dendritic cells include lymphoid dendritic cells of the spleen, Langerhans cells of the epidermis, and veiled cells in the blood circulation.

Collectively, dendritic cells are classified as a group based on their morphology, high levels of surface MHC-class II expression, and absence of certain other surface markers expressed on T cells, B cells, monocytes, and natural killer cells. In particular, monocyte-derived dendritic cells (also referred to as monocytic dendritic cells) usually express CD11c, CD80, CD86, and are HLA-DR<sup>+</sup>, but are CD14<sup>-</sup>.

In contrast, monocytic dendritic cell precursors (typically monocytes) are usually CD14<sup>+</sup>. Monocytic dendritic cell precursors can be obtained from any tissue where they reside, particularly lymphoid tissues such as the spleen, bone marrow, lymph nodes and thymus. Monocytic dendritic cell precursors also can be isolated from the circulatory system. Peripheral blood is a readily accessible source of monocytic dendritic cell precursors.

Umbilical cord blood is another source of monocytic dendritic cell precursors. Monocytic dendritic cell precursors can be isolated from a variety of organisms in which an immune response can be elicited. Such organisms include animals, for example, including humans, and non-human animals, such as, primates, mammals (including dogs, cats, mice, and rats), birds (including chickens), as well as transgenic species thereof.

In certain embodiments, the monocytic dendritic cell precursors and/or immature dendritic cells can be isolated from a healthy subject or from a subject in need of immunostimulation, such as, for example, a cancer patient or other subject for whom cellular immunostimulation can be beneficial or desired (*i.e.*, a subject having a bacterial or viral infection, and the like). Dendritic cell precursors and/or immature dendritic cells also can be obtained from an HLA-matched healthy individual for administration to an HLA-matched subject in need of immunostimulation.



### Dendritic Cell Precursors and Immature Dendritic Cells

Methods for isolating cell populations enriched for dendritic cell precursors and immature dendritic cells from various sources, including blood and bone marrow, are known in the art. For example, dendritic cell precursors and immature dendritic cells can be isolated by collecting heparinized blood, by apheresis or leukapheresis, by preparation of buffy coats, rosetting, centrifugation, density gradient centrifugation (*e.g.*, using Ficoll® (such as FICOLL-PAQUE®), PERCOLL® (colloidal silica particles (15-30 nm diameter) coated with non-dialyzable polyvinylpyrrolidone (PVP)), sucrose, and the like), differential lysis of cells, filtration, and the like. In certain embodiments, a leukocyte population can be prepared, such as, for example, by collecting blood from a subject, defibrinating to remove the platelets and lysing the red blood cells. Dendritic cell precursors and immature dendritic cells can optionally be enriched for monocytic dendritic cell precursors by, for example, centrifugation through a PERCOLL® gradient, antibody panning, and the like.

Dendritic cell precursors and immature dendritic cells optionally can be prepared in a closed, aseptic system. As used herein, the terms "closed, aseptic system" or "closed system" refer to a system in which exposure to non-sterile, ambient, or circulating air or other non-sterile conditions is minimized or eliminated. Closed systems for isolating dendritic cell precursors and immature dendritic cells generally exclude density gradient centrifugation in open top tubes, open air transfer of cells, culture of cells in tissue culture plates or unsealed flasks, and the like. In a typical embodiment, the closed system allows aseptic transfer of the dendritic cell precursors and immature dendritic cells from an initial collection vessel to a sealable tissue culture vessel without exposure to non-sterile air.

Another reported method for isolating dendritic cell precursors is to use a commercially treated plastic substrate (*e.g.*, beads or magnetic beads) to selectively remove adherent monocytes and other "non-dendritic cell precursors." (See, *e.g.*, U.S. Patent Nos. 5,994,126 and 5,851,756.) The adherent monocytes and non-dendritic cell precursors are discarded while the non-adherent cells are retained for *ex vivo* culture and maturation. In another method, apheresis cells were cultured in plastic culture bags to which plastic, *i.e.*, polystyrene or styrene, microcarrier beads were added to increase the surface area of the bag. The cells were cultured for a sufficient period of time for cells to adhere to the beads and the non-adherent cells were washed from the bag. (Maffei, *et al.*, *Transfusion* 40:1419-1420 (2000); WO 02/44338, incorporated herein by reference).

In certain other embodiments, monocytic dendritic cell precursors are isolated by adherence to a monocyte-binding substrate, as disclosed in U.S. Patent Application No.

PCT/US02/23865, filed July 25, 2002, the disclosure of which is incorporated by reference herein. For example, a population of leukocytes (*e.g.*, isolated by leukapheresis) can be contacted with a monocytic dendritic cell precursor adhering substrate. When the population of leukocytes is contacted with the substrate, the monocytic dendritic cell precursors in the leukocyte population preferentially adhere to the substrate. Other leukocytes (including other potential dendritic cell precursors) exhibit reduced binding affinity to the substrate, thereby allowing the monocytic dendritic cell precursors to be preferentially enriched on the surface of the substrate.

Suitable substrates include, for example, those having a large surface area to volume ratio. The substrate can be, for example, a particulate or fibrous substrate. Suitable particulate substrates include, for example, glass particles, plastic particles, glass-coated plastic particles, glass-coated polystyrene particles, and other beads suitable for protein absorption. Fibrous substrates suitable for use in the present invention include microcapillary tubes and microvillous membrane and the like. The particulate or fibrous substrate usually allows the adhered monocytic dendritic cell precursors to be eluted without substantially reducing the viability of the adhered cells. A particulate or fibrous substrate can be substantially non-porous to facilitate elution of monocytic dendritic cell precursors or dendritic cells from the substrate. A "substantially non-porous" substrate is a substrate in which at least a majority of pores present in the substrate are smaller than the cells to minimize entrapping cells in the substrate.

Adherence of the monocytic dendritic cell precursors to the substrate can optionally be enhanced by the addition of a binding media. Suitable binding media include monocytic dendritic cell precursor culture media (*e.g.*, AIM-V<sup>®</sup>, RPMI 1640, DMEM, X-VIVO 15<sup>®</sup>, and the like) supplemented, individually or in any combination, with for example, cytokines (*e.g.*, Granulocyte/Macrophage Colony Stimulating Factor (GM-CSF), Interleukin 4 (IL-4), or Interleukin 13 (IL-13)), blood plasma, serum (*e.g.*, human serum, such as autologous or allogenic sera), purified proteins, such as serum albumin, divalent cations (*e.g.*, calcium and/or magnesium ions) and other molecules that aid in the specific adherence of monocytic dendritic cell precursors to the substrate, or that prevent adherence of non-monocytic dendritic cell precursors to the substrate. In certain embodiments, the blood plasma or serum can be heated-inactivated. The heat-inactivated plasma can be autologous or heterologous to the leukocytes.

Following adherence of monocytic dendritic cell precursors to the substrate, the non-adhering leukocytes are separated from the monocytic dendritic cell

precursor/substrate complexes. Any suitable means can be used to separate the non-adhering cells from the complexes. For example, the mixture of the non-adhering leukocytes and the complexes can be allowed to settle, and the non-adhering leukocytes and media decanted or drained. Alternatively, the mixture can be centrifuged, and the supernatant containing the non-adhering leukocytes decanted or drained from the pelleted complexes.

In another method, monocytic dendritic cell precursors can be isolated from a cell population enriched in leukocytes prepared by the use of a tangential flow filtration device. A tangential flow filtration device useful for the isolation of a cell population enriched in monocytic dendritic cell precursors can comprise a remover unit having a cross-flow chamber, a filtrate chamber and a filter disposed therebetween. The filter is in fluid communication on one side, the retentate surface, with the cross-flow chamber, and on the other side, the filtrate surface, with the filtrate chamber. The cross-flow chamber has an inlet adapted to introduce a sample of blood constituents comprising leukocytes into the cross-flow chamber and parallel to the retentate surface of the filter. An outlet is also provided in the cross-flow chamber centrally disposed in a portion of the chamber opposite the retentate surface of the filter. The filter suitable for use in the tangential flow filtration device typically has an average pore size ranging from about 1 to about 10 microns. The filter can have an average pore size of about 3 to about 7 microns. A means for providing a predetermined input rate of the sample into the inlet of the cross-flow chamber and a means for controlling a filtration rate of filtrate through the filter and into the filtrate chamber can also be included. The filtration rate controlling means limits the rate of filtration to less than the unopposed filtration rate for the filter. The sample comprising blood constituents can be provided by a source device such as a leukapheresis device or a container comprising a sample collected from a leukapheresis device.

Monocytic dendritic cell precursors and cell populations enriched for the precursors can be cultured *ex vivo* for differentiation, and partial maturation and/or expansion. (As used herein, isolated immature dendritic cells, dendritic cell precursors, and other cells, refers to cells that, by human hand, exist apart from their native environment, and are therefore not a product of nature. Isolated cells can exist in purified form, in semi-purified form, or in a non-native environment.) Briefly, *ex vivo* differentiation typically involves culturing monocytic dendritic cell precursors, or populations of cells having dendritic cell precursors, in the presence of one or more differentiation agents. Suitable differentiating agents can include, for example, cellular growth factors (*e.g.*, cytokines such as (GM-CSF), Interleukin 4 (IL-4), Interleukin 7 (IL-7), Interleukin 13 (IL-13), and/or

combinations thereof). In certain embodiments, the monocytic dendritic cells precursors are differentiated to form monocyte-derived immature dendritic cells.

The dendritic cell precursors can be cultured and differentiated in suitable culture conditions. Suitable tissue culture media include, but are not limited to, AIM-V<sup>®</sup>, RPMI 1640, DMEM, X-VIVO 15<sup>®</sup>, and the like. The tissue culture media can be supplemented with serum, amino acids, vitamins, cytokines, such as GM-CSF and/or IL-4, IL-7, IL-13, divalent cations, and the like, to promote differentiation of the cells. In certain embodiments, the dendritic cell precursors can be cultured in serum-free media. The culture conditions can optionally exclude any animal-derived products. A typical cytokine combination used with dendritic cell culture medium comprises about 500 units/ml each of GM-CSF and IL-4, IL-7, or IL-13. Dendritic cell precursors, when differentiated to form immature dendritic cells, are phenotypically similar to skin Langerhans cells. Immature dendritic cells typically are CD14<sup>-</sup> and CD11c<sup>+</sup>, express low levels of CD86 and CD83, and are able to capture soluble antigens via specialized endocytosis.

Dendritic cell maturation agents can include, for example, but are not limited to, BCG, IFN $\gamma$ , LPS, TNF $\alpha$ , and the like. Effective amounts of BCG typically range from about 10<sup>5</sup> to 10<sup>7</sup> cfu per milliliter of tissue culture media. Effective amounts of IFN $\gamma$  typically range from about 100-1000 U per milliliter of tissue culture media. Bacillus Calmette-Guerin (BCG) is an avirulent strain of *M. bovis*. As used herein, BCG refers to whole BCG as well as cell wall constituents, BCG-derived lipoarabidomannans, and other BCG components. BCG is optionally inactivated, such as heat-inactivated BCG, formalin-treated BCG, and the like.

The immature DCs are typically contacted with effective amounts of BCG and IFN $\gamma$  for about one hour to about 10 hours. The immature dendritic cells can be cultured and partially matured in suitable maturation culture conditions. Suitable tissue culture media include AIM-V<sup>®</sup>, RPMI 1640, DMEM, X-VIVO 15<sup>®</sup>, and the like. The tissue culture media can be supplemented with amino acids, vitamins, cytokines, such as GM-CSF and/or IL-15, divalent cations, and the like, to promote maturation of the cells. A typical cytokine combination is about 500 units/ml of GM-CSF and 100 ng/ml of IL-15.

Partial maturation of immature dendritic cells can be monitored by methods known in the art for dendritic cells. Cell surface markers can be detected in assays familiar to the art, such as flow cytometry, immunohistochemistry, and the like. The cells can also be monitored for cytokine production (e.g., by ELISA, another immune assay, or by use of an

oligonucleotide array). In DCs cultured and partially matured according to the present invention in the presence of a dendritic cell maturation agent, such as GM-CSF and IL-4, the levels of phosphorylated JAK2 (janus activated kinase 2) can be measured to indicate the initiation of maturation by methods well known in the art. The induction of the expression of cell surface markers and cytokines, as well as the phosphorylation of signaling molecules, *e.g.*, jak2, is also known as an indicator that dendritic cells are conditioned for induction of an immune response once the cells have been administered to an individual.

The immature dendritic cells are subject to maturation only for a time period to partially mature the dendritic cells. Fully mature DCs lose the ability to take up antigen and display up-regulated expression of costimulatory cell surface molecules and various cytokines. Specifically, mature DCs express higher levels of MHC class I and II antigens than immature dendritic cells, and mature dendritic cells are generally identified as being CD80<sup>+</sup>, CD83<sup>+</sup>, CD86<sup>+</sup>, and CD14<sup>-</sup>. Greater MHC expression leads to an increase in antigen density on the DC surface, while up regulation of costimulatory molecules CD80 and CD86 strengthens the T cell activation signal through the counterparts of the costimulatory molecules, such as CD28 on the T cells.

Fully mature dendritic cells are not preferred for the present invention because they do not efficiently process antigen. Further, immature dendritic cells as used in prior methods are not desired because the immunosuppressive environment typically found within a tumor, including substantial concentrations of cytokines known to prevent the processing of antigen by immature dendritic cells. In the present invention, partial maturation of the immature dendritic cells down regulates cytokine receptors on the surface of the cell making them less sensitive or responsive to any immunosuppressive effects of cytokines present in the intratumoral space and provides for cells that can efficiently process tumor antigens present within the intratumoral space. Once the partially mature dendritic cells have matured within the tumor as measured by the production of, for example, TNF- $\alpha$  and IL-12, and the expression of the chemokine receptor CCR7, the cells can migrate to the lymph nodes where the cells will contact T cells to up regulate the immune response to the tumor antigens.

According to yet another aspect of the invention, DCs can be preserved, *e.g.*, by cryopreservation either before maturation or following partial maturation prior to administration to a patient. Cryopreservation agents which can be used include but are not limited to dimethyl sulfoxide (DMSO), glycerol, polyvinylpyrrolidone, polyethylene glycol, albumin, dextran, sucrose, ethylene glycol, i-erythritol, D-ribitol, D-mannitol, D-sorbitol, i-inositol, D-lactose, choline chloride, amino acids, methanol, acetamide, glycerol

monoacetate, and inorganic salts. A controlled slow cooling rate can be critical. Different cryoprotective agents and different cell types typically have different optimal cooling rates. The heat of fusion phase where water turns to ice typically should be minimal. The cooling procedure can be carried out by use of, *e.g.*, a programmable freezing device or a methanol bath procedure. Programmable freezing apparatuses allow determination of optimal cooling rates and facilitate standard reproducible cooling. Programmable controlled-rate freezers such as Cryomed or Planar permit tuning of the freezing regimen to the desired cooling rate curve.

After thorough freezing, monocytic precursor cells, immature DCs or partially mature DCs can be rapidly transferred to a long-term cryogenic storage vessel. In a typical embodiment, samples can be cryogenically stored in liquid nitrogen ( $-196^{\circ}\text{C}$ .) or its vapor ( $-165^{\circ}\text{C}$ .). Considerations and procedures for the manipulation, cryopreservation, and long term storage of hematopoietic stem cells, particularly from bone marrow or peripheral blood, is largely applicable to the cells of the invention. Such a discussion can be found, for example, in the following references, incorporated by reference herein: Taylor *et al.*, *Cryobiology* 27:269-78 (1990); Gorin, *Clinics in Haematology* 15:19-48 (1986); Bone-Marrow Conservation, Culture and Transplantation, Proceedings of a Panel, Moscow, Jul. 22-26, 1968, International Atomic Energy Agency, Vienna, pp. 107-186.

Frozen cells are preferably thawed quickly (*e.g.*, in a water bath maintained at  $37^{\circ}\text{C}$ - $41^{\circ}\text{C}$ ) and chilled immediately upon thawing. It may be desirable to treat the cells in order to prevent cellular clumping upon thawing. To prevent clumping, various procedures can be used, including but not limited to the addition before and/or after freezing of DNase (Spitzer *et al.*, *Cancer* 45: 3075-85 (1980)), low molecular weight dextran and citrate, hydroxyethyl starch (Stiff *et al.*, *Cryobiology* 20: 17-24 (1983)), and the like. The cryoprotective agent, if toxic in humans, should be removed prior to therapeutic use of the thawed partially matured DCs. One way in which to remove the cryoprotective agent is by dilution to an insignificant concentration. Once frozen partially matured DCs have been thawed and recovered, they can be administered as described herein with respect to non-frozen partially matured DCs.

#### In vivo Administration of Cell Populations

Methods are provided for administration of partially mature dendritic cells, or a cell population enriched and containing such cells, to a subject having a tumor. In certain

### *In vivo* Administration of Cell Populations

Methods are provided for administration of partially mature dendritic cells, or a cell population enriched and containing such cells, to a subject having a tumor. In certain embodiments, such methods are performed by obtaining dendritic cell precursors or immature dendritic cells, differentiating and partially maturing those cells in the presence of a dendritic cell maturation agent, such as BCG and IFN $\gamma$ . The partially matured dendritic cells can be formulated with physiologically acceptable carriers, excipients, buffers and/or diluents using methods and compositions well known to the skilled artisan. The partially mature dendritic cells can be administered directly to a subject in need of immunostimulation. Typically, about  $10^2$  to about  $10^{10}$  cells are suspended in a pharmaceutically acceptable carrier. The cells are injected either into the tumor directly or into a region near to, adjacent to, or in circulatory or lymphatic contact with the tumor to ensure that the cells have access to the tumor antigens. For example, but not by limitation, the cells can be administered directly into a tumor, into the tumor bed subsequent to surgical removal or resection of the tumor, peritumorally, into a draining lymph node in direct contact with the tumor, into a blood vessel or lymph duct leading into, or feeding a tumor or organ afflicted by the tumor, *e.g.*, the portal vein or a pulmonary vein or artery, and the like. The administration of the partially mature dendritic cells of the invention can be either simultaneous with or subsequent to other treatments for the tumor, such as chemotherapy or radiation therapy. In addition, the dendritic cells can also be formulated or compounded into a slow release matrix for implantation into a region in or around the tumor or tumor bed such that cells are slowly released into the tumor, or tumor bed, for contact with the tumor antigens.

Partially mature dendritic cells of the present invention can be administered by any means appropriate for the formulation and mode of administration. For example, the cells can be combined with a pharmaceutically acceptable carrier and administered with a syringe, a catheter, a cannula, and the like. As above, the cells can be formulated in a slow release matrix. When administered in this fashion, the formulation can be administered by a means appropriate for the matrix used. Other methods and modes of administration applicable to the present invention are well known to the skilled artisan.

Compositions of the present invention can be used by themselves in the treatment of an individual. In addition, the compositions can be used in combination with any other method to treat a tumor. For example, the methods of the present invention can be used in combination with surgical resection of a tumor, chemotherapy (cytotoxic drugs,

apoptotic agents, antibodies, and the like), radiation therapy, immune therapy (administration of antigen specific mature activated dendritic cells, NK cells, antibodies specific for tumor antigens, *etc.*), and the like. Any and all of these methods can also be used in any combination. Combination treatments can be concurrent or sequential and can be administered in any order as determined by the treating physician.

In another embodiment, the dendritic cells and the recipient subject have the same MHC (HLA) haplotype. Methods of determining the HLA haplotype of a subject are known in the art. In a related embodiment, the partially mature dendritic cells are allogenic to the recipient subject. The allogenic cells are typically matched for at least one MHC allele (*e.g.*, sharing at least one but not all MHC alleles). In a less typical embodiment, the dendritic cells and the recipient subject are all allogeneic with respect to each other, but all have at least one MHC allele in common.

### Examples

In the present example, monocytic dendritic cell precursors, isolated by tangential flow filtration, were cultured at a concentration of  $10^6$  monocytes/ml in X-VIVO 15<sup>®</sup> supplemented with 2% human serum albumin (HSA) and 500 U/ml of GM-CSF in either flasks or cell bags. After 5 days, the DCs from the flasks and the bags were either left untreated (iDCs) or were matured for 4 hours in the presence of BCG (1:400 dilution) and interferon  $\gamma$  (IFN $\gamma$ ; 500 U/ml) for four hours (pmDCs). The cells were collected, washed and mixed with an equal number of fluorescently labeled irradiated PKH67-A549 tumor cells for 48 hours. Following the incubation, the cells were washed, stained with a phycoerythrin (PE) labeled monoclonal antibody specific for CD11c, and analyzed by flow cytometry. The values provided in Table 1 represent the percentage of CD11c<sup>+</sup> cells within the DC gate that were also positive for labeled PKH67 antigen (MFI = mean fluorescent intensity). Controls included DCs mixed with PKH67-A549 cells just prior to analysis.

Treatment	% CD11c <sup>+</sup> PKH67 <sup>+</sup> Cells	MFI
iDCs-flask	87.6	235.2
pmDCs-flask	90.7	610.2
iDCs-bag	72.8	403.5
pmDCs-bag	91.3	557.5



These data demonstrate that the partially matured DCs were better at antigen uptake in this assay than immature dendritic cells, as measured by either the percentage of cells that take up antigen or by the amount of material picked up by the tumor cells.

In another example, a leukapheresis was carried out on a patient subsequent to surgery for removal of a tumor growth. Monocytes were purified from the leukapheresis product by tangential flow filtration. The purified monocytes were differentiated as described above and partially matured by exposure to heat killed and formalin fixed BCG ( $5 \times 10^6$  cfu/ml) IFN $\gamma$  (500 U/ml). The partially matured dendritic cells were formulated for administration and injected into the tumor bed of the individual, under ultrasound guidance. Dendritic cells isolated from the individual were found to be capable of inducing a tumor specific cytotoxic T cell response when measured *in vitro*. Further, it was found that the time to tumor recurrence was prevented or substantially delayed.

The previous examples are provided to illustrate, but not to limit, the scope of the claimed inventions. Other variants of the inventions will be readily apparent to those of ordinary skill in the art and encompassed by the appended claims. All publications, patents, patent applications and other references cited herein and are also incorporated by reference herein in their entirety.

WHAT IS CLAIMED IS:

1                   1.       A method for producing an anti-tumor immune response comprising  
2 administration to an individual with a tumor a cell population comprising dendritic cells that  
3 have been partially matured *in vitro*, wherein the partially matured dendritic cells can take up  
4 and process antigen and are enabled to induce an anti-tumor immune response subsequent to  
5 administration to the individual.

1                   2.       The method according to claim 1, wherein the dendritic cells are  
2 obtained from skin, spleen , bone marrow, thymus, lymph nodes, umbilical cord blood, or  
3 peripheral blood.

1                   3.       The method according to claim 2, wherein the dendritic cells are  
2 obtained from the individual to be treated.

1                   4.       The method according to claim 2, wherein the dendritic cells are  
2 obtained from a healthy individual HLA-matched to the individual to be treated.

1                   5.       The method according to claim 1, wherein the dendritic cells are  
2 partially matured in the presence of a dendritic cell maturation agent.

1                   6.       The method according to claim 5, wherein the dendritic cell maturation  
2 agent is Bacillus Calmette-Guerin (BCG), interferon  $\gamma$  (IFN $\gamma$ ), lipopolysaccharide (LPS),  
3 tumor necrosis factor  $\alpha$  (TNF $\alpha$ ), or a combination thereof.

1                   7.       The method according to claim 6, wherein BCG comprises whole  
2 BCG, cell wall constituents of BCG, BCG-derived lipoarabidomannans, or BCG components.

1                   8.       The method according to claim 6, wherein the BCG is heat-inactivated  
2 BCG or formalin-treated BCG.

1                   9.       The method according to claim 6, wherein the effective amount of  
2 BCG is about  $10^5$  to  $10^7$  cfu per milliliter of tissue culture media and the effective amount of  
3 IFN $\gamma$  is about 100 to about 1000 Units per milliliter of tissue culture media.

1                   10.      The method according to claim 1, wherein the partially matured  
2 dendritic cells are administered directly into the tumor.

1                    13.     The method according to claim 1, wherein the partially matured  
2     dendritic cells are administered into a lymph node directly draining a tumor area.

1                    15.     The method according to claim 1, wherein the partially matured  
2     dendritic cells are administered into the circulatory system such that the cells are delivered to  
3     the tumor or tumor afflicted organ.

1           17.     The method according to claim 16, wherein the partially matured  
2     dendritic cells are administered prior to, simultaneous with, or subsequent to radiation  
3     therapy, chemotherapy, or combinations thereof.

1                    19.     The composition according to claim 18, wherein the partially mature  
2     dendritic cells do not express the cell surface marker CD83.

20. The composition according to claim 18, wherein the composition comprises about  $10^2$  to about  $10^{10}$  partially matured dendritic cells.

1                   21.     The composition according to claim 18, wherein the partially matured  
2 dendritic cells have been cryopreserved subsequent to partial maturation.

1                   22.     The composition according to claim 18, wherein the partially matured  
2 dendritic cells have been isolated from a patient to whom they are to be administered.

1                   23.     The composition according to claim 18, wherein the partially matured  
2 dendritic cells have been HLA matched to an individual to whom they are to be administered.

1                   24.     The composition according to claim 18, wherein the partially matured  
2 dendritic cells are administered directly into the tumor.

1                   25.     The composition according to claim 18, wherein the partially matured  
2 dendritic cells are administered into a tumor bed subsequent to surgical removal or resection  
3 of the tumor.

1                   26.     The composition according to claim 18, wherein the partially matured  
2 dendritic cells are administered to an tissue area surrounding the tumor.

1                   27.     The composition according to claim 18, wherein the partially matured  
2 dendritic cells are administered into a lymph node directly draining a tumor area.

1                   28.     The composition according to claim 18, wherein partially matured  
2 dendritic cells are administered directly to a circulatory vessel duct that delivers blood or  
3 lymph to the tumor or a tumor afflicted organ.

1                   29.     The method according to claim 18, wherein the partially matured  
2 dendritic cells are administered into the circulatory system such that the cells are delivered to  
3 the tumor or tumor afflicted organ.

PATENT

Attorney Docket No.: 020093-004000

**ADMINISTRATION OF DENDRITIC CELLS PARTIALLY MATURED  
*IN VITRO* FOR THE TREATMENT OF TUMORS**

**ABSTRACT OF THE DISCLOSURE**

The present invention provides populations of cells comprising partially matured dendritic cells that can be used for administration individuals having a tumor. Partially matured dendritic cells, those contacted with a dendritic cell maturation agent for 1 to 10 hours, efficiently take up and process tumor antigens in area of the tumor site, complete maturation and subsequently migrate to the lymph nodes of a treated individual. Once in the lymph node the now fully mature antigen presenting dendritic cells secrete the appropriate cytokines (*e.g.*,  $\text{TNF}\alpha$  and IL-12) and contact T cells inducing a powerful anti-tumor immune response.

SE 5014672 v1

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**